

Contents lists available at ScienceDirect

Journal of Cardiology Cases

journal homepage: www.elsevier.com/locate/jccase

Case Report

Late perforation of anterior mitral leaflet after surgical resection of the subaortic membrane

Hatem Ari (MD)^{a,*}, Fatih Kahraman (MD)^a, Akif Arslan (MD)^a, Kadir Ceviker (MD)^b, Fatih Aksoy (MD)^a^a Department of Cardiology, Suleyman Demirel University School of Medicine, Isparta, Turkey^b Department of Cardiovascular Surgery, Suleyman Demirel University School of Medicine, Isparta, Turkey

ARTICLE INFO

Article history:

Received 26 May 2015

Received in revised form 28 July 2015

Accepted 13 August 2015

Keywords:

Mitral leaflet perforation

Mitral regurgitation

Endothelial injury

ABSTRACT

A 54-year-old woman who underwent surgical resection of the subaortic membrane 10 years earlier presented with new onset dyspnea. Cardiovascular examination revealed 3–4/6 pansystolic murmur at the apex. She was found to have severe mitral regurgitation (MR) with transthoracic echocardiography; 2D and real-time-3D transesophageal echocardiography demonstrated severe MR through anterior mitral leaflet perforation with precise localization. The patient was treated with surgery in which the perforated segment was closed by direct suture technique and discharged on postoperative 5th day.

<Learning objective: Late anterior mitral leaflet perforation after surgical or interventional procedures has rarely been reported. We present this case to emphasize the role of traumatic injury to weak endothelial surfaces such as a valve leaflet in the development of late leaflet perforation after surgical or interventional procedures.>

© 2015 Japanese College of Cardiology. Published by Elsevier Ltd. All rights reserved.

Introduction

Mitral leaflet perforation (MLP) is a rarely encountered clinical condition that may result in acute severe mitral regurgitation (MR). MLP may be found together with mitral valve aneurysm (MVA) or in isolated form. Anterior MLP with or without aneurysm is more frequently observed compared to posterior MLP and is commonly encountered in the early or late course of aortic valve infective endocarditis (IE) [1]. Primary pathology or degeneration of valvular tissue is other mechanistic cause of MLP other than IE [2,3]. MLP cases have been rarely reported during or subsequent to different surgical or interventional cardiac procedures that are relevant to mitral valve and left ventricular outflow tract (LVOT) [4–10].



In this context, we present a case of late small anterior mitral leaflet aneurysm perforation 10 years after surgical resection of the subaortic membrane without any evidence of previous or active endocarditis.

Case report

A 54-year-old woman who had a history of open-heart surgery for subaortic membrane resection 10 years earlier was admitted to

our clinic with complaints of new onset dyspnea and palpitation. Cardiovascular examination revealed 3–4/6 pansystolic murmur at the apex. Electrocardiography showed atrial fibrillation with a rapid ventricular rate. Transthoracic echocardiography (TTE) showed severe eccentric regurgitant jet from the LVOT to the left atrium through the perforated segment of the basal part of anterior mitral leaflet just below the aortic non-coronary cusp (Fig. 1A; supplementary material online, Video 1). No residual subaortic membrane was seen. Moderate aortic regurgitation and tricuspid fibrocalcific aortic valve were also seen with TTE imaging. The patient was on warfarin therapy due to atrial fibrillation and no mitral valve regurgitation was noted in the regular follow-up visits. Physical examination, laboratory markers, and blood cultures were negative for IE. Transesophageal echocardiography (TEE) revealed echo dropout at the top of a small aneurysmal sac which was located at the basal part of the A2 scallop of the anterior mitral leaflet (Fig. 1B; supplementary material online, Video 2). Real-time three-dimensional TEE (RT-3D-TEE) imaging clearly showed systolic explication of a small aneurysm with perforation (Fig. 2A) at the region of the basal part of the A2 scallop of the anterior mitral leaflet and diastolic collapse of the perforated aneurysmal sac (Fig. 2B). The patient was referred to surgery for mitral valve repair after coronary angiography, which was completely normal. Intraoperative inspection verified echocardiography by revealing the small aneurysm with perforation at the A2 scallop of the anterior mitral leaflet, 2 mm next to mitral

* Corresponding author at: Araştırma ve Uygulama Hastanesi, Süleyman Demirel Üniversitesi, 32260 Isparta, Turkey. Tel.: +90 2462112000; fax: +90 2462112830. E-mail address: hatem_ari@hotmail.com (H. Ari).



annulus. The perforated segment was closed by direct suture technique and the patient was discharged on postoperative 5th day.

Discussion

The most common form of subaortic stenosis is the membranous type and the membrane may be attached to the anterior leaflet of the mitral valve and encircle the LVOT. The location of this membrane ranges from just below the aortic valve to further down into the left ventricle. Many patients present with associated aortic regurgitation (AR). Definitive therapy for subaortic membrane is generally surgical resection.

Although mitral leaflet tears may be seen as an acute complication during surgical subaortic membrane resection [4], cases of anterior mitral leaflet perforation have been reported in the relatively late or late postoperative period after surgical resection of the subaortic membrane [5,6]. Ozyuksel et al. reported a case similar to ours [5]. In this case, anterior mitral leaflet perforation and severe MR associated with the perforation had developed 4 months after the surgical treatment of discrete subaortic membrane. These authors suggested inappropriate surgical technique as the reason for perforation. Anterior mitral leaflet perforation has been reported even 30 years after surgical subaortic membrane resection [6].

Fibrous continuity between the aortic annuli and anterior mitral leaflet where the subaortic membrane might be attached or continues to this location could be injured with an apparent tear,

perforation, or be minimally damaged such as by endothelial denudation during surgery. Continuing tissue reaction after the index tissue injury and endothelial denudation could have led to late perforation with or without the formation of aneurysms by weakening the mitral leaflet. Systolic left ventricular pressure stress and diastolic hemodynamic stress due to moderate aortic regurgitation, which is frequently observed in these patients, to the injured leaflet tissue may be contributory factors by weakening the leaflet tissue leading to aneurysm and subsequent perforation. It is possible that index trauma during the initial surgery along with other continual contributing factors mentioned above could have resulted in the observations of this case.

MLP with or without aneurysm has rarely been reported during or subsequent to several surgical or interventional cardiac procedures that are relevant to the mitral valve and LVOT [7–10]. All of these interventional procedures potentially have minimal or more traumatic contact to the mitral leaflet suggesting the role of index traumatic injury to the valve leaflet for the development of late perforation.

We present this report to underline the importance of trauma to the weak endothelial surface such as the valve leaflet, in the surgical or interventional procedures, even if it is minimal.

Conflict of interest

The authors report no financial relationships or conflicts of interest regarding the content herein.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.jccase.2015.08.010>.

References

- [1] Guler A, Karabay CY, Gursoy OM, Guler Y, Candan O, Akgun T, Bulut M, Pala S, Izgi AI, Esen AM, Kirma C, Ozkan M. Clinical and echocardiographic evaluation of mitral valve aneurysms: a retrospective, single center study. *Int J Cardiovasc Imaging* 2014;30:535–41.
- [2] Tewari S, Moorthy N, Sinha N. Ruptured mitral valve aneurysm: an uncommon cause of acute dyspnoea. *Echocardiography* 2010;27:E119–21.
- [3] Kim DJ, Cho KI, Jun HJ, Kim YJ, Song YJ, Jhi JH, Chon MG, Kim SM, Lee HG, Kim TI. Perforated mitral valve aneurysm in the posterior leaflet without infective endocarditis. *J Cardiovasc Ultrasound* 2012;20:100–2.
- [4] Rattes MF, Sochowski RA, Baird M, Chan KL. Intraoperative transesophageal echocardiographic demonstration of mitral leaflet tear following resection of a subaortic membrane. *Can J Cardiol* 1992;8:785–7.
- [5] Ozyuksel A, Yildirim O, Onsel I, Bilal MS. Severe mitral regurgitation due to anterior mitral leaflet perforation after surgical treatment of discrete subaortic stenosis. *BMJ Case Rep* 2014;2014. <http://dx.doi.org/10.1136/bcr-2014-204463>. pii: bcr2014204463.
- [6] Bustamante J, Flórez S, García-Morán E, Bustamante E, Gómez-Martínez Mde L, Tamayo E. Mitral insufficiency 30 years after resection of subvalvular aortic stenosis. *Asian Cardiovasc Thorac Ann* 2010;18:202.
- [7] Cozzarin A, Cianciulli TF, Guidoin R, Zhang Z, Lax JA, Saccheri MC, García Escudero A, Estrada JE. Core valve prosthesis causes anterior mitral leaflet perforation resulting in severe mitral regurgitation. *Can J Cardiol* 2014;30:1108.e11–3.
- [8] Javed U, Smith TW, Rogers JH. Percutaneous repair of anterior mitral leaflet perforation. *J Invasive Cardiol* 2012;24:134–7.
- [9] Ohtsuka T, Murakawa Y, Hayami N, Kotsuka Y, Takamoto S. Surgically repaired delayed mitral regurgitation after radiofrequency catheter ablation for Wolff-Parkinson-White syndrome. *Pacing Clin Electrophysiol* 2002;25:1142–3.
- [10] Fisch-Thomsen M, Jensen JK, Egeblad H, Wierup P, Nørgaard BL. Mitral valve perforation appearing years after radiofrequency ablation. *J Heart Valve Dis* 2011;20:351–2.